

## A note on convex renorming and fragmentability

A K MIRMOSTAFAEE

Department of Mathematics, Damghan University of Sciences, P.O. 36715/364,  
Damghan, Iran  
E-mail: kmirmostafae@du.ac.ir

MS received 29 November 2004

**Abstract.** Using the game approach to fragmentability, we give new and simpler proofs of the following known results: (a) If the Banach space admits an equivalent Kadec norm, then its weak topology is fragmented by a metric which is stronger than the norm topology. (b) If the Banach space admits an equivalent rotund norm, then its weak topology is fragmented by a metric. (c) If the Banach space is weakly locally uniformly rotund, then its weak topology is fragmented by a metric which is stronger than the norm topology.

**Keywords.** Fragmentability of Banach spaces; topological games; renorming of Banach spaces.

### 1. Introduction

Let  $(X, \tau)$  be a topological space and  $\rho$  be a metric on  $X$ . Given  $\varepsilon > 0$ , a nonempty subset  $A$  of  $X$  is said to be *fragmented by  $\rho$  down to  $\varepsilon$*  if each nonempty subset of  $A$  contains a nonempty  $\tau$ -relatively open subset of  $\rho$ -diameter less than  $\varepsilon$ .  $A$  is called *fragmented by  $\rho$*  if  $A$  is fragmented by  $\rho$  down to  $\varepsilon$  for each  $\varepsilon > 0$ . The set  $A$  is said to be  *$\sigma$ -fragmented by  $\rho$*  if for every  $\varepsilon > 0$ ,  $A$  can be expressed as  $A = \bigcup_{n=1}^{\infty} A_{n,\varepsilon}$  with each  $A_{n,\varepsilon}$  fragmented by  $\rho$  down to  $\varepsilon$ .

The notion of fragmentability was originally introduced in [3] to investigate the existence of nice selections for upper semicontinuous compact-valued mappings. The notion of  $\sigma$ -fragmentability appeared in [1] in order to study Banach spaces, the weak topology of which is  $\sigma$ -fragmented by the norm (such Banach spaces are said to be  $\sigma$ -fragmentable). Since then, these two concepts have been playing an important role in the study of the geometry of Banach spaces.

Kenderov and Moors [4] used the following topological game to characterize fragmentability of a topological space  $X$ : Two players  $\Sigma$  and  $\Omega$  alternatively select subsets of  $X$ .  $\Sigma$  starts the game by choosing some nonempty subset  $A_1$  of  $X$ . Then  $\Omega$  chooses some nonempty relatively open subset  $B_1$  of  $A_1$ . In general, if the selection  $B_n \neq \emptyset$  of the player  $\Omega$  is already specified, the player  $\Sigma$  makes the next move by selecting an arbitrary nonempty set  $A_{n+1}$  contained in  $B_n$ . Continuing the game the two players generate a sequence of sets

$$A_1 \supset B_1 \supset \cdots \supset A_n \supset B_n \supset \cdots$$

which is called a play and is denoted by  $p = (A_i, B_i)_{i=1}^{\infty}$ . If

$$p_1 = (A_1), \dots, p_n = (A_1, B_1, \dots, A_n)$$

are the first 'n' move of some play (of the game), then  $p_n$  is called the  $n$ th *partial play* of the game. The player  $\Omega$  is said to have won the play  $p$  if  $\bigcap_{i=1}^{\infty} A_i = \bigcap_{i=1}^{\infty} B_i$  contains at most one point. Otherwise the player  $\Sigma$  is said to be the winner in this play. Under the term *strategy*  $s$  for  $\Omega$ -player, we mean a rule by means of which the player  $\Omega$  makes his/her choices. More precisely, the strategy  $s$  is a sequence of mappings  $s = \{s_n\}_{n \geq 1}$ , which are defined inductively as follows:  $s_1$  assigns to each possible first move  $A_1$  of  $\Sigma$ -player a nonempty relatively open subset  $B_1 = s_1(A_1)$ . Therefore, the domain of  $s_1$  is the set of all nonempty subsets of  $X$  and  $s_1$  assigns to each such an element a nonempty relatively open subset of it. The domain of  $s_2$  consists of triples of the type  $(A_1, B_1, A_2)$ , where  $A_1$  is from the domain of  $s_1$ ,  $B_1 = s_1(A_1)$  and  $A_2$  is an arbitrary nonempty subset of  $B_1$ .  $s_2$  assigns to such a triple a nonempty relatively open subset  $B_2 = s_2(A_1, B_1, A_2)$  of  $A_2$ . In general, the domain of  $s_{n+1}$  consists of partial plays of the type

$$(A_1, \dots, A_i, B_i, A_{i+1}, \dots, A_{n+1}),$$

where, for every  $i \leq n$ ,  $(A_1, \dots, A_i)$  is from the domain of  $s_i$ ,  $B_i = s_i(A_1, \dots, A_i)$  and  $A_{n+1}$  is an arbitrary nonempty subset of  $B_n$ . To every element from its domain  $s_{n+1}$  assigns a nonempty relatively open subset  $B_{n+1}$  of  $A_{n+1}$ .

A play  $p = (A_i, B_i)_{i \geq 1}$  is called an  $s$ -play if  $B_i = s_i(p_i)$  for each  $i \geq 1$ .  $s$  is called a *winning strategy* for the player  $\Omega$  if he/she wins every  $s$ -play. If the space  $X$  is fragmentable by a metric  $d(\cdot, \cdot)$ , then  $\Omega$  has an obvious winning strategy  $s$ . Indeed, to each partial play  $p_n$  this strategy puts into correspondence some nonempty subset  $B_n \subset A_n$  which is relatively open in  $A_n$  and has  $d$ -diameter less than  $1/n$ . Clearly, the set  $\bigcap_{i \geq 1} A_i = \bigcap_{i \geq 1} B_i$  has at most one point because it has zero  $d$ -diameter. Kenderov and Moors have shown that the existence of a winning strategy for the player  $\Omega$  characterizes fragmentability, that is,

**Theorem 1.1 [4].** *The topological space  $X$  is fragmentable if and only if the player  $\Omega$  has a winning strategy.*

Of special interest is the case when the topology generated by the fragmenting metric contains the original topology of the space (in this case it is said that  $X$  is *fragmented by a metric which is stronger than its topology*).

**Theorem 1.2 [4].** *The topological space  $X$  is fragmentable by a metric stronger than its topology if and only if the player  $\Omega$  has a strategy  $a$  such that, for every  $s$ -play  $p = (A_i, B_i)_{i \geq 1}$  the intersection  $\bigcap_{i=1}^{\infty} A_i = \bigcap_{i=1}^{\infty} B_i$  is either empty or contains just one point  $x_0$  and for every neighborhood  $U$  of  $x_0$  there exists some  $k$  such that  $A_i \subset U$  for all  $i > k$ .*

This characterization of fragmentability has some applications (see e.g. [4-6]). In [5], it is shown that fragmentability and  $\sigma$ -fragmentability of the weak topology in a Banach space are related to each other in the following way:

**Theorem 1.3 ([5], Theorems 1.3, 1.4 and 2.1).** *For a Banach space  $X$  the following are equivalent:*

- (i)  $(X, \text{weak})$  is  $\sigma$ -fragmented by the norm (i.e.  $X$  is  $\sigma$ -fragmented);
- (ii)  $(X, \text{weak})$  is fragmented by a metric which is stronger than the weak topology;
- (iii)  $(X, \text{weak})$  is fragmented by a metric which is stronger than the norm topology;
- (iv) There exists a strategy  $s$  for the player  $\Omega$  in  $(X, \text{weak})$  such that, for every  $s$ -play  $p = (A_i, B_i)_{i \geq 1}$  either  $\bigcap_{i \geq 1} B_i = \emptyset$  or  $\lim_{i \rightarrow \infty} \text{norm-diam}(B_i) = 0$ .

- (v) *There exists a strategy  $s$  for the player  $\Omega$  in  $(X, \text{weak})$  such that, for every  $s$ -play  $p = (A_i, B_i)_{i \geq 1}$  either  $\bigcap_{i \geq 1} B_i = \emptyset$  or every sequence  $\{x_i\}_{i \geq 1}$  with  $x_i \in B_i, i \geq 1$  has a weak cluster point.*

Moreover, we have the following: The norm  $\|\cdot\|$  of a Banach space  $X$  is said to be *Kadec* if the norm topology and the weak topology coincide on the unit sphere  $\{x \in X: \|x\| = 1\}$ . In [2], it was shown that every Banach space with Kadec norm is  $\sigma$ -fragmented. It follows that there exists a strategy for the player  $\Omega$  satisfying condition (iv) from the theorem of Kenderov and Moors. In the next section, we will directly construct such a strategy (without using the theorem of Kenderov and Moors).

The norm  $\|\cdot\|$  of a Banach space  $X$  is said to be *rotund* (or *strictly convex*) if the unit sphere  $\{x \in X: \|x\| = 1\}$  does not contain nontrivial line segments. Ribarska has shown in [7] that the weak topology of a rotund Banach space is fragmented by a metric. By the abovementioned characterization of fragmentability it follows that the player  $\Omega$  has a winning strategy. In the next section we will directly define such a strategy (without using the result of Ribarska and the mentioned theorem of Kenderov and Moors). Moreover, if the norm of  $X$  is weakly locally uniformly rotund, then the strategy we construct satisfies condition (v) from the above theorem of Kenderov and Moors. Recall that the Banach space  $X$  is called *locally uniformly rotund* (resp. *weakly locally uniformly rotund*) if  $\lim_{n \rightarrow \infty} \|x_n - x\| = 0$  (resp.  $\text{weak-lim}(x_n - x) = 0$ , whenever  $\lim_{n \rightarrow \infty} \|(x_n + x)/2\| = \lim_{n \rightarrow \infty} \|x_n\| = \|x\|$ ).

## 2. Description of the strategies

*Lemma 1. Let  $X$  be a Banach space with Kadec norm. Then, for every  $\varepsilon > 0$  and  $x \in X$ , there exists some positive number  $\alpha_{\varepsilon, x}$  and a weakly open set  $W_{\varepsilon, x} \ni x$  such that  $\|y - x\| < \varepsilon$  whenever  $y \in W_{\varepsilon, x}$  and  $|\|y\| - \|x\|| \leq \alpha_{\varepsilon, x}$ .*

*Proof.* If  $x = 0$ , it suffices to put  $W_{\varepsilon, x} = X$  and to take as  $\alpha_{\varepsilon, x}$  any positive number smaller than  $\varepsilon/2$ . Suppose  $x \neq 0$  and take a convex weakly open neighborhood  $G$  of  $x$  such that the norm diameter of  $G \cap \{z: \|z\| = \|x\|\}$  is less than  $\varepsilon/2$ . Define  $\alpha_{\varepsilon, x} > 0$  to be smaller than  $\varepsilon/2, \|x\|$  and such that  $\alpha_{\varepsilon, x}B \subset (G - x)/2$  (as usual  $B$  stands for the closed unit ball of  $X$ ). Put  $W_{\varepsilon, x} := x + (G - x)/2 = (x + G)/2$ . Let  $y \in W_{\varepsilon, x}$  and  $|\|y\| - \|x\|| < \alpha_{\varepsilon, x}$ . Then we have

$$\begin{aligned} (\|x\|/\|y\|)y &= ((\|x\|/\|y\|)y - y) + y = (\|x\| - \|y\|)y/\|y\| + y \\ &\in \|\|y\| - \|x\|\|B + W_{\varepsilon, x} \subset \alpha_{\varepsilon, x}B + W_{\varepsilon, x} \subset (G - x)/2 + (G + x)/2 = G. \end{aligned}$$

Hence  $\|(\|x\|/\|y\|)y - x\| < \varepsilon/2$ . Finally we have

$$\|y - x\| \leq \|y - (\|x\|/\|y\|)y\| + \|(\|x\|/\|y\|)y - x\| < \alpha_{\varepsilon, x} + \varepsilon/2 < \varepsilon. \quad \square$$

We also need the following result:

*Lemma 2. ([5], Proposition 2.1). If the closed unit ball  $B$  of a Banach space  $X$  admits a strategy  $s$  with the property (iv) of Theorem 1.3, then the whole space also admits such a strategy.*

**Theorem 2.1.** *Let  $X$  be a Banach space with Kadec norm. Then there exists a strategy  $s$  for the player  $\Omega$  in  $(B, \text{weak})$  such that, for every  $s$ -play  $p = (A_i, B_i)_{i \geq 1}$  either  $\bigcap_{i \geq 1} B_i = \emptyset$  or  $\lim_{i \rightarrow \infty} \text{norm-diam}(B_i) = 0$ .*

*Proof.* Let  $\|\cdot\|$  denote the Kadec norm on  $X$  and  $A_1$  be the first choice of  $\Sigma$ -player. By Lemma 2, we may assume that  $A_1 \subset B$ , where  $B$  denotes the closed unit ball of  $X$ . Put

$$\rho_1 = \sup\{\|x\| : x \in A_1\} \quad \text{and} \quad \varepsilon_1 = 1.$$

Two cases may happen.

- (1) There is an element  $x_1 \in A_1$  such that  $\alpha_{\varepsilon_1, x_1} + \|x_1\| > \rho_1$ . Then we take such a point  $x_1$  and define  $s_1(A_1) = B_1 := W_{\varepsilon_1, x_1} \cap A_1 \setminus (\|x_1\| - \alpha_{\varepsilon_1, x_1})B$  and  $\varepsilon_2 := \varepsilon_1/2$ . Then for each  $y \in B_1$ ,  $\|y\| \leq \rho_1 < \alpha_{\varepsilon_1, x_1} + \|x_1\|$  and  $\|y\| \geq \|x_1\| - \alpha_{\varepsilon_1, x_1}$ . Therefore, by Lemma 1,  $\|y - x_1\| < \varepsilon_1$ . Hence  $\| \cdot \| - \text{diam}(B_1) < 2\varepsilon_1$ .
- (2) For every  $x \in A_1$ ,  $\alpha_{\varepsilon_1, x} + \|x\| \leq \rho_1$ . Then,

$$s_1(A_1) = B_1 := A_1 \setminus (1/2)\rho_1 B$$

and set  $\varepsilon_2 = \varepsilon_1$ . Suppose the mappings  $(s_i)_{i \leq n}$  participating in the definition of a strategy for player  $\Omega$  have already been defined. Let  $(A_i, B_i)_{1 \leq i \leq n}$  be a partial play which is generated by the strategy mappings defined so far. This partial play is accompanied by the numbers  $\{\varepsilon_i\}_{1 \leq i \leq n}$  and the points  $x_1, \dots, x_n$ . If  $A_{n+1}$  is the next move of the player  $\Sigma$ , we put

$$\rho_{n+1} = \sup\{\|x\| : x \in A_{n+1}\}$$

and consider the following two possible cases:

- (1) There exists an element  $x_{n+1} \in A_{n+1}$ , such that  $\alpha_{\varepsilon_{n+1}, x_{n+1}} + \|x_{n+1}\| > \rho_{n+1}$ . In this case, we take such a point  $x_{n+1}$ , define

$$s_{n+1}(A_1, \dots, A_{n+1}) = B_{n+1} := W_{\varepsilon_{n+1}, x_{n+1}} \cap A_{n+1} \setminus (\|x_{n+1}\| - \alpha_{\varepsilon_{n+1}, x_{n+1}})B$$

and set  $\varepsilon_{n+2} = \varepsilon_{n+1}/2$ . As above one shows that in this case  $\| \cdot \| - \text{diam}(B_{n+1}) < 2\varepsilon_{n+1}$ .

- (2) For every point  $x \in A_{n+1}$ ,  $\alpha_{\varepsilon_{n+1}, x} + \|x\| \leq \rho_{n+1}$ . In this case, we define

$$s_{n+1}(A_1, \dots, A_{n+1}) = B_{n+1} := A_{n+1} \setminus \left(1 - \frac{1}{(n+2)}\right) \rho_{n+1} B$$

and set  $\varepsilon_{n+2} = \varepsilon_{n+1}$ . In this way the strategy  $s = (s_i)_{i \geq 1}$  for the  $\Omega$ -player is already defined.

Suppose  $(A_i, B_i)_{i \geq 1}$  is an  $s$ -play with  $x \in \bigcap_{n \geq 1} A_n$  and  $\lim_{n \rightarrow \infty} \| \cdot \| - \text{diam}(B_n) \neq 0$ . Then there exists some  $\delta > 0$ , such that  $\| \cdot \| - \text{diam}(B_n) > \delta$  for each  $n \in \mathbb{N}$ . This means that for all but finitely many  $n$ , the case (2) happens and thus  $\{\varepsilon_n\}$  is eventually constant:  $\varepsilon_n = \varepsilon > 0$  for all  $n > k$ . Since  $x \in \bigcap_{n \geq 1} A_n$ ,

$$\left(1 - \frac{1}{n}\right) \rho_n < \|x\| < \rho_n, \quad \text{for all } n.$$

Let  $\rho_n \searrow \rho$ . Then the above inequality shows that  $\|x\| = \rho$ . On the other hand,  $\alpha_{\varepsilon, x} + \|x\| = \alpha_{\varepsilon_n, x} + \|x\| \leq \rho_n$  for  $n > k$  which implies the contradiction  $\alpha_{\varepsilon, x} + \|x\| = \|x\|$ .  $\square$

*Remark 2.2.* Lemma 1 directly implies that Banach spaces with Kadec norm are  $\sigma$ -fragmentable. Actually, Theorem 2.3 of [2] indirectly implies that every Kadec renormable Banach space  $X$  has a countable cover by sets of small local norm diameter, i.e., for each  $\varepsilon > 0$ , it is possible to write  $X = \bigcup_{n \in \mathbb{N}} X_{n,\varepsilon}$  such that for each  $n \in \mathbb{N}$  and  $x \in X_{n,\varepsilon}$ , there exists an open neighborhood  $V_x$  of  $x$  such that the norm diameter of  $V_x \cap X_{n,\varepsilon}$  is less than  $\varepsilon$ . Using Lemma 1, we can give another proof of this result.

**PROPOSITION 2.3.**

*Let  $X$  be a Banach space with Kadec norm. Then for every  $\varepsilon > 0$  there exists a countable cover of  $X$ ,  $X = \bigcup_{i \geq 0} X_i$ , such that, for every  $x \in X_i$ , there exists a weakly open neighborhood  $W$  of  $x$  such that  $W \cap X_i$  is contained in  $x + \varepsilon B$ , in particular the points of  $X_i$  have weak neighborhoods with norm-diameter smaller than  $2\varepsilon$ .*

*Proof.* Given  $\varepsilon > 0$  consider, for  $k = 1, 2, \dots$ , and  $n = 0, 1, 2, \dots$ , the sets  $X_{kn} = \{x \in X : \alpha_{\varepsilon,x} > 2/k, \text{ and } n/k \leq \|x\| \leq (n+1)/k\}$ . Clearly,  $X$  is covered by  $X_{kn}$ . Put  $W := W_{\varepsilon,x}$ . By Lemma 1 the set  $W \cap X_{kn}$  is contained in  $x + \varepsilon B$ .  $\square$

**Theorem 2.4.** *Let  $X$  be a Banach space.*

- (a) *If the norm of  $X$  is rotund, then  $(X, \text{weak})$  is fragmentable by a metric.*
- (b) *If the norm of  $X$  is weakly locally uniformly rotund, then  $(X, \text{weak})$  is fragmented by a metric which is stronger than the norm topology.*

*Proof.* According to Theorems 1.2 and 1.3 and Lemma 2, it is enough to show that in  $(B, \text{weak})$  the player  $\Omega$  has a winning strategy  $s$  such that, for every  $s$ -play  $p = (A_i, B_i)_{i \geq 1}, \bigcap_{i \geq 1} B_i$  has at most one point and in case (b) either  $\bigcap_{i \geq 1} B_i = \emptyset$  or every sequence  $\{y_n\}, y_n \in B_n, n \geq 1$  is weakly convergent to the element of  $\bigcap_{i \geq 1} B_i$ . Let  $\|\cdot\|$  be the equivalent norm on  $X$  and  $\Sigma$  start a game by choosing a nonempty subset  $A_1$  of  $B$ . Define

$$\rho_1 = \sup\{\|x\| : x \in A_1\}.$$

Choose an element  $x_1 \in A_1$  such that  $\|x_1\| > \rho_1 - 1/2$  and find some  $\mu_1 \in X^*$  such that  $\|\mu_1\| = 1$  and  $\mu_1(x_1) = \|x_1\|$ . Define

$$s_1(A_1) = B_1 := \{x \in A_1 : \mu_1(x) > \rho_1 - 1/2\}$$

as the first choice of  $\Omega$ -player. Then for each  $x \in B_1$ , we have

$$\rho_1 - 1/2 < \mu_1(x) \leq \|x\| \leq \rho_1.$$

Suppose that the finite sequence  $\{x_k\}_{k \leq n}$  of points of  $X$ ,  $\{\mu_k\}_{k \leq n}$  of elements of  $X^*$ , and the partial play  $p_n = (A_1, \dots, B_n)$  have already been specified so that for each  $x \in B_k, k \leq n$  the inequality

$$\rho_k - \frac{1}{k+1} < \mu_k(x) < \|x\| \leq \rho_k$$

holds. Let  $A_{n+1}$  be the answer of  $\Sigma$ -player to  $p_n$ . Put

$$\rho_{n+1} = \sup\{\|x\| : x \in A_{n+1}\}$$

and find some  $x_{n+1} \in A_{n+1}$ ,  $\|x_{n+1}\| > \rho_{n+1} - \frac{1}{n+2}$ . Take some  $\mu_{n+1} \in X^*$ ,  $\|\mu_{n+1}\| = 1$  with  $\mu_{n+1}(x_{n+1}) = \|x_{n+1}\|$  and define

$$s_{n+1}(A_1, \dots, A_{n+1}) = B_{n+1} = \left\{ x \in A_{n+1} : \mu_{n+1}(x) > \rho_{n+1} - \frac{1}{n+2} \right\},$$

as the next choice of the player  $\Omega$ . Clearly, for each  $x \in B_{n+1}$ , the inequality

$$\rho_{n+1} - \frac{1}{n+2} < \mu_{n+1}(x) < \|x\| \leq \rho_{n+1}$$

holds. Thus, by induction on  $n$ , we have shown that the  $\Omega$ -player can choose sets of the form

$$B_n = \left\{ x \in A_n : \mu_n(x) > \rho_n - \frac{1}{n+1} \right\},$$

where  $\|\mu_n\| = 1$  and  $\rho_n = \sup\{\|x\| : x \in A_n\}$  for each  $n \in N$ .

Let  $\cap_{n \geq 1} B_n \neq \emptyset$  and  $\mu$  be a weak\* cluster point of  $\{\mu_n\}$ . Then for each  $x \in \cap_{n \geq 1} B_n$ , the inequality

$$\rho_n - \frac{1}{n+1} < \mu_n(x) < \|x\| \leq \rho_n$$

for each  $n \in N$  implies that  $\mu(x) = \|x\| = \rho$ , where  $\rho$  is the limit of the decreasing sequence  $\{\rho_n\}$ . It follows that for each  $x, y \in \cap_{n \geq 1} B_n$ , we have  $\mu(x) = \|x\| = \|y\| = \mu(y)$ . Rotundity of  $X$  implies that  $x = y$ , thus, in this case,  $\cap_{n \geq 1} B_n$  has at most one point. In case (b), suppose that  $x \in \cap_{n \geq 1} B_n$ . If  $y_n \in B_n$ , the inequality

$$\rho_n - \frac{1}{n+1} < \frac{1}{2} \mu_n(x + y_n) \leq \frac{1}{2} \|x + y_n\| \leq \frac{1}{2} (\|x\| + \|y_n\|) \leq \rho_n$$

shows that  $\lim_{n \rightarrow \infty} \|(x + y_n)/2\| = \lim_{n \rightarrow \infty} \|y_n\| = \|x\| = \rho$ . Since  $(X, \|\cdot\|)$  is weakly locally uniformly rotund, it follows that  $\lim_{n \rightarrow \infty} (x - y_n) = 0$ . By Theorem 1.2, the space is fragmented by a metric stronger than the weak topology. This completes the proof.  $\square$

**Remark 2.5.** It is well-known that locally uniformly rotund norms are Kadec. Therefore statement (b) from the above theorem follows from Theorem 2.1 as well.

## Acknowledgement

The author is indebted to P Kenderov for the useful remarks while this work was in progress.

## References

- [1] Jayne J E, Namioka I and Rogers C A, Topological properties of Banach spaces, *Proc. London Math. Soc.* **(3)66** (1993) 651–672
- [2] Jayne J E, Namioka I and Rogers C A,  $\sigma$ -Fragmentable Banach spaces I, *Mathematika* **39** (1992) 161–188

- [3] Jayne J E and Rogers C A, Borel selectors for upper semi-continuous set-valued maps, *Acta Math.* **56** (1985) 41–79
- [4] Kenderov P S and Moors W B, Game characterization of fragmentability of topological spaces, *Mathematics and Education in Mathematics* (1996) pp. 8–18 (Proceedings of the 25th Spring Conference of the Union of Bulgarian Mathematicians, April 1996, Kazanlak, Bulgaria)
- [5] Kenderov P S and Moors W B, Fragmentability and sigma-fragmentability of Banach spaces, *J. London Math. Soc.* **(2)60** (1999) 203–223
- [6] Mirmostafaei A K, On non-fragmentability of Banach spaces, *Proc. Indian Acad. Sci. (Math. Sci.)* **108(2)** (1998) 163–167
- [7] Ribarska N K, A note on fragmentability of some topological spaces, *C. R. I'Acad, Bulgare des Sci.* t. 43, No. 7 (1990) 13–15